

Children's Acetaminophen (Tylenol)

| Weight | 80mg dropper, Infant Tylenol | Elixer 160mg/5ml Children's Tylenol | Chewable Tablets 80mg | Junior Strength 160mg | Suppository 80,220,325,650mg |
|-----------|------------------------------|-------------------------------------|-----------------------|-----------------------|------------------------------|
| 6-11 lbs | ½ dropper, 0.4ml | ¼ tsp | N/A | N/A | 80mg, ½ sup |
| 12-17 lbs | 1 dropper, 0.8ml | ½ tsp | 1 tablet | N/A | 80mg, 1 sup |
| 18-23 lbs | 1 ½ droppers, 1.2ml | ¾ tsp | 1 ½ tablets | N/A | 120mg, 1 sup |
| 24-35 lbs | 2 droppers, 1.6ml | 1 tsp | 2 tablets | N/A | 325mg, ½ sup |
| 36-47 lbs | N/A | 1 ½ tsp | 3 tablets | N/A | 325mg, ¾ sup |
| 48-59 lbs | N/A | 2 tsp | 4 tablets | 2 tablets | 325mg, 1 sup |
| 60-71 lbs | N/A | 2 ½ tsp | 5 tablets | 2 ½ tablets | 325mg, 1 sup |
| 72-95 lbs | N/A | 3 tsp | 6 tablets | 3 tablets | 650mg, ¾ sup |
| 96+ lbs | N/A | N/A | N/A | 4 tablets | 650mg, 1 sup |

Do not give Tylenol more often than every 4 hours or more than 5 doses per day.

Note: Since Tylenol pediatric products are available without prescription, parents are warned on the package label to consult a physician for the use by children under 2 yrs or for the use longer than 5 days. Contact your physician immediately in the case of accidental overdose

Claritin Liquid

2-6 years: 1 tsp daily, do not take more than 1 tsp in 24 hours

6+ years: 2 tsp daily, do not take more than 2 tsp in 24 hours

Children's Zyrtec Liquid

6-24 months: ½ tsp at bedtime

2-6 years: ½ to 1 tsp daily, maximum dose is ½ tsp every 12 hours or 1 tsp in 24 hours

6 + years: 1-2 tsp daily depending on severity of symptoms, do not exceed 2 tsp in 24 hours

Children's Benadryl *Not Recommended until 12 months of age*

| Weight | 12.5 mg = 1 tsp | 25 mg capsule |
|-----------|-----------------|-----------------|
| 6-11 lbs | Not recommended | Not recommended |
| 12-17 lbs | ½ tsp | Not recommended |
| 18-23 lbs | 1 tsp | Not recommended |
| 24-35 lbs | 1 ½ tsp | Not recommended |
| 36-47 lbs | 2 tsp | 1 capsule |
| 48-59 lbs | 3 tsp | 1 capsule |
| 60-71 lbs | 3-4 tsp | 2 capsules |
| 72-95 lbs | 4 tsp | 2 capsules |
| 96+ lbs | 4 tsp | 2 capsules |

Children's Ibuprofen (same as Motrin or Advil) *Not recommended until 6 months of age*

| Age | Weight | Infant Drops 50mg/1.25ml | Ibuprofen Liquid 100mg/5ml | Ibuprofen Chewable 50mg | Ibuprofen Chewable 100mg |
|--------------|-----------|-----------------------------|----------------------------------|-------------------------------|--------------------------------|
| 6-11 months | 13-17 lbs | 1.25ml | ½ tsp | 1 tablet | ½ tablet |
| 12-23 months | 18-23 lbs | 1.875ml | ¾ tsp | 1 ½ tablets | ¾ tablet |
| 2-3 yrs | 24-35 lbs | N/A | 1 tsp | 2 tablets | 1 tablet |
| 4-5 yrs | 36-47 lbs | N/A | 1 ½ tsp | 2 ½ tablets | 1 ½ tablets |
| 6-8 yrs | 48-59 lbs | N/A | 2 tsp | 4 tablets | 2 tablets |
| 9-10 yrs | 60-71 lbs | N/A | 2 ½ tsp | 5 tablets | 2 ½ tablets |
| 11-12 yrs | 72-95 lbs | N/A | 3 tsp | 6 tablets | 3 tablets |

One dose may be given every 6-8 hours as needed for fever control. Do not exceed 4 doses in 24 hours.